

$\angle \theta$  &  $(x, y)$  be  
Cartesian coordinates  $\frac{\partial z}{\partial \theta} = \frac{\partial z}{\partial x} \cdot \frac{\partial x}{\partial \theta} + \frac{\partial z}{\partial y} \cdot \frac{\partial y}{\partial \theta} = \frac{\partial z}{\partial x} (-r \sin \theta) + \frac{\partial z}{\partial y} (r \cos \theta)$   
in the plane and

let  $(r, \theta)$  be polar  $\frac{\partial(\frac{\partial z}{\partial \theta})}{\partial \theta} = \frac{\partial}{\partial \theta} \left( \frac{\partial z}{\partial x} \cdot \frac{\partial x}{\partial \theta} \right) + \frac{\partial}{\partial \theta} \left( \frac{\partial z}{\partial y} \cdot \frac{\partial y}{\partial \theta} \right)$   
coordinates.

Express  $\frac{\partial^2 z}{\partial \theta^2}$  =  $\boxed{① \frac{\partial}{\partial \theta} \left( \frac{\partial z}{\partial x} \right)} \cdot \frac{\partial x}{\partial \theta} + \frac{\partial}{\partial \theta} \left( \frac{\partial z}{\partial y} \right) \cdot \frac{\partial y}{\partial \theta}$   
in Cartesian  
coordinates.

$$+ \boxed{② \frac{\partial}{\partial \theta} \left( \frac{\partial z}{\partial y} \right)} \frac{\partial y}{\partial \theta} + \frac{\partial}{\partial \theta} \left( \frac{\partial z}{\partial x} \right) \cdot \frac{\partial z}{\partial y}$$

①

$$\begin{aligned} \frac{\partial}{\partial \theta} \left( \frac{\partial z}{\partial x} \right) &= \frac{\partial \left( \frac{\partial z}{\partial x} \right)}{\partial x} \cdot \frac{\partial x}{\partial \theta} + \frac{\partial \left( \frac{\partial z}{\partial x} \right)}{\partial y} \cdot \frac{\partial y}{\partial \theta} \\ &= \frac{\partial^2 z}{\partial x^2} (-r \sin \theta) + \frac{\partial^2 z}{\partial y^2} \cdot (r \cos \theta). \end{aligned}$$

②

$$\begin{aligned} \frac{\partial}{\partial \theta} \left( \frac{\partial z}{\partial y} \right) &= \frac{\partial \left( \frac{\partial z}{\partial y} \right)}{\partial x} \cdot \frac{\partial x}{\partial \theta} + \frac{\partial \left( \frac{\partial z}{\partial y} \right)}{\partial y} \cdot \frac{\partial y}{\partial \theta} \\ &= \frac{\partial^2 z}{\partial x \partial y} \cdot (-r \sin \theta) + \frac{\partial^2 z}{\partial y^2} \cdot (r \cos \theta) \end{aligned}$$

$$\begin{aligned} \frac{\partial}{\partial \theta} \left( \frac{\partial z}{\partial \theta} \right) &= ① \cdot (-r \sin \theta) + (-r \sin \theta) \cdot \frac{\partial z}{\partial x} + \\ &\quad ② \cdot (r \cos \theta) + (r \cos \theta) \cdot \frac{\partial z}{\partial y} \end{aligned}$$

I will leave the rest to  
you guys ...

Taylor Series with Another Base Point

$$f(x,y) = f(a,b) + f_x(a,b)(x-a) + f_y(a,b)(y-b) \\ + \frac{1}{2!} [f_{xx}(a,b)(x-a)^2 + 2f_{xy}(a,b)(x-a)(y-b) + f_{yy}(y-b)^2] + \dots$$

→ Linear approx. involves  $f_x, f_y$

→ Quadratic approx. involves  $f_{xx}, f_{xy}, f_{yy}$

Example 2: We find the 2<sup>nd</sup> order Taylor expansion of

$f(x,y) = \sqrt{1+4x^2+y^2}$  about (1,2) and use it to compute approximately  $f(1.1, 2.05)$ .

$$f(x,y) = \sqrt{1+4x^2+y^2} \quad f(1,2) = 3$$

$$f_x(x,y) = \frac{4x}{\sqrt{1+4x^2+y^2}} \quad f_x(1,2) = \frac{4}{3}$$

$$f_y(x,y) = \frac{y}{\sqrt{1+4x^2+y^2}} \quad f_y(1,2) = \frac{2}{3}$$

$$f_{xx}(x,y) = \frac{4}{\sqrt{1+4x^2+y^2}} - \frac{16x^2}{\sqrt{1+4x^2+y^2}} \quad f_{xx}(1,2) = \frac{20}{27}$$

$$f_{xy}(1,2) = -\frac{8}{27}$$

$$f_{yy}(1,2) = \frac{5}{27}$$

$$f(x,y) = 3 + \frac{4}{3}(x-1) + \frac{2}{3}(y-2) + \frac{10}{27}(x-1)^2 - \frac{8}{27}(x-1)(y-2) \\ + \frac{5}{27}(y-2)^2 + \text{Error}$$

$$f(1.1, 2.05) \approx 3 + \frac{4}{3}(1-1) + \frac{2}{3}(0.05) + \frac{10}{27}(0.01) \\ - \frac{8}{27}(0.005) + \frac{5}{27}(0.0025) = 3.1691.$$

Example 3. Find and classify all critical points.

$$\sin(x) + \sin(y) + \sin(x+y) \text{ on } (0, \pi) \times (0, \pi)$$

$$f_x = \cos(x) + \cos(x+y)$$

$$f_y = \cos(y) + \cos(x+y)$$

$$f_x = 0, f_y = 0$$

$$\begin{cases} \cos x + \cos(x+y) = 0 \Rightarrow \cos x = -\cos(x+y) \\ \cos y + \cos(x+y) = 0 \Rightarrow \cos y = -\cos(x+y) \end{cases}$$

$$\therefore \cos y = \cos x$$

$$\therefore y = x \quad 0 < x, y < \pi$$

$$\cos(x) + \cos(2x) = 0$$

$$\cos(x) + 2\cos^2 x - 1 = 0$$

$$\cos(x) = \frac{-1 \pm \sqrt{1+8}}{4}$$

$$\cos(x) = -1 \quad \text{or} \quad \cos(x) = \frac{1}{2}$$

↑  
not in the  
range

Solution

$$x = \frac{\pi}{3}$$

$$CP: \left(\frac{\pi}{3}, \frac{\pi}{3}\right)$$

$$Hf = \begin{vmatrix} \frac{\partial^2 f}{\partial x^2} & \frac{\partial^2 f}{\partial x \partial y} \\ \frac{\partial^2 f}{\partial y \partial x} & \frac{\partial^2 f}{\partial y^2} \end{vmatrix} = \begin{vmatrix} -\sin x - \sin(x+y) & -\sin(x+y) \\ -\sin(x+y) & -\sin y - \sin(x+y) \end{vmatrix}$$

$$Hf\left(\frac{\pi}{3}, \frac{\pi}{3}\right) = \begin{vmatrix} -ns & -\frac{n^2}{2} \\ -\frac{\sqrt{3}}{2} & -\sqrt{3} \end{vmatrix}$$

$$\det(A_1) = -\sqrt{3}$$

$$\det(A_2) = \frac{9}{4} > 0$$

- + - + ... local max.